


U.S. Fish and Wildlife Service

# Coastal Wetland Trends in the Narragansett Bay Estuary During the 20th Century: Rhode Island



# Coastal Wetland Trends in the Narragansett Bay Estuary During the 20th Century: Rhode Island

November 2004

A National Wetlands Inventory Cooperative Interagency Report

# Coastal Wetland Trends in the Narragansett Bay Estuary During the 20th Century: Rhode Island

Ralph W. Tiner<sup>1</sup>, Irene J. Huber<sup>2</sup>, Todd Nuerminger<sup>2</sup>, and Aimée L. Mandeville<sup>3</sup>

<sup>1</sup>U.S. Fish & Wildlife Service  
National Wetlands Inventory Program  
Northeast Region  
300 Westgate Center Drive  
Hadley, MA 01035

<sup>2</sup>Natural Resources Assessment Group  
Department of Plant and Soil Sciences  
University of Massachusetts  
Stockbridge Hall  
Amherst, MA 01003

<sup>3</sup>Department of Natural Resources Science  
Environmental Data Center  
University of Rhode Island  
1 Greenhouse Road, Room 105  
Kingston, RI 02881

November 2004

National Wetlands Inventory Cooperative Interagency Report between  
U.S. Fish & Wildlife Service, University of Massachusetts-Amherst, University of Rhode  
Island, and Rhode Island Department of Environmental Management

This report should be cited as: Tiner, R.W., I.J. Huber, T. Nuerminger, and A.L. Mandeville. 2004. Coastal Wetland Trends in the Narragansett Bay Estuary During the 20th Century: Rhode Island. U.S. Fish and Wildlife Service, Northeast Region, Hadley, MA. In cooperation with the University of Massachusetts-Amherst and the University of Rhode Island. National Wetlands Inventory Cooperative Interagency Report. 37 pp. plus appendices.

## Table of Contents

	Page
Introduction	1
Study Area	1
Methods	5
Data Compilation	5
Geospatial Database Construction and GIS Analysis	8
Results	9
Baywide 1996 Status	9
Coastal Wetlands and Waters	9
500-foot Buffer Zone	9
Baywide Trends 1951/2 to 1996	15
Coastal Wetland Trends	15
500-foot Buffer Zone Around Coastal Wetlands	15
Trends for Pilot Study Areas	25
Conclusions	35
Acknowledgments	36
References	37
Appendices	
A. Baywide Summary Tables for the Narragansett Bay Estuary	
B. Summary Tables for Individual Study Areas	

## List of Tables

No.	Page
1. Aerial photography used for this study.	6
2. Causes of wetland losses, gains, and changes in type.	7
3. 1996 status of coastal wetlands and waters in the Narragansett bay Estuary.	10
4. Extent of altered coastal wetlands for the Narragansett Bay Estuary in 1996.	13
5. Land use/cover in the 500-foot buffer around coastal wetlands in the Narragansett Bay Estuary in 1996.	14
6. Trends in coastal wetlands and waters in the Narragansett Bay Estuary from the 1950s to the 1990s.	16
7. Nature and causes of coastal wetland changes in the Narragansett Bay Estuary from the 1950s to the 1990s.	19
8. Land use/cover changes in the 500-foot buffer surrounding tidal wetlands in the Narragansett Bay Estuary from the 1950s to the 1990s.	23
9. Status and trends in coastal wetlands for specific study areas.	27
10. Nature and causes of coastal wetland and deepwater habitat trends for Allins Cove.	29
11. Nature and causes of coastal wetland and deepwater habitat trends for Calf Pasture Point.	30
12. Nature and causes of coastal wetland and deepwater habitat trends for Jacobs Point.	31
13. Nature and causes of coastal wetland and deepwater habitat trends for Palmer River.	32
14. Nature and causes of coastal wetland and deepwater habitat trends for Sachuset Point.	33
15. Nature and causes of coastal wetland and deepwater habitat trends for Wesquage Pond.	34

### Tables in Appendix A:.

- 1-A. Changes in estuarine emergent wetlands in the Narragansett Bay Estuary: 1950s to 1990s.
- 2-A. Changes in estuarine scrub-shrub wetlands in the Narragansett Bay Estuary: 1950s to 1990s.
- 3-A. Changes in estuarine unconsolidated shores in the Narragansett Bay Estuary: 1950s to 1990s.
- 4-A. Changes in vegetated coastal wetlands in the Narragansett Bay Estuary: 1950s to 1990s.
- 5-A. Changes in nonvegetated coastal wetlands in the Narragansett Bay Estuary: 1950s to 1990s.

## **List of Tables** (continued)

### Tables in Appendix B:

- 1-B. Trends in estuarine wetlands for Allins Cove from the 1930s to the 1950s and from the 1950s to the 1990s.
- 2-B. Trends in estuarine wetlands for Calf Pasture Point from the 1930s to the 1950s and from the 1950s to the 1990s.
- 3-B. Trends in estuarine wetlands for Jacobs Point from the 1930s to the 1950s and from the 1950s to the 1990s.
- 4-B. Trends in estuarine wetlands for Palmer River from the 1930s to the 1950s and from the 1950s to the 1990s.
- 5-B. Trends in estuarine wetlands for Sachuest Point from the 1930s to the 1950s and from the 1950s to the 1990s.
- 6-B. Trends in estuarine wetlands for Wesquage Pond from the 1930s to the 1950s and from the 1950s to the 1990s.

## **List of Figures**

No.	Page
1. Location of the Narragansett Bay Estuary and its drainage area.	2
2. Limits of the Narragansett Bay Estuary as defined for this study.	3
3. Location of six pilot areas within the Narragansett Bay Estuary.	4
4. Percent loss of estuarine emergent wetland in the Narragansett Bay Estuary.	21
5. Percent gain in estuarine emergent wetland in the Narragansett Bay Estuary.	21
6. Percent change in estuarine scrub-shrub wetland in the Narragansett Bay Estuary.	22
7. Percent change in estuarine unconsolidated shore in the Narragansett Bay Estuary.	22

## **Introduction**

The Rhode Island Department of Environmental Management's Narragansett Bay Estuary Program's (NBEP) goal is to protect and preserve Narragansett Bay through conserving and restoring natural resources and enhancing water quality. NBEP accomplishes this through a variety of projects, including interagency partnerships and community involvement. To manage these valuable resources, NBEP wanted baseline information on coastal wetlands and their buffers. With the aid of the University of Massachusetts (UMass), University of Rhode Island (URI), and the U.S. Fish and Wildlife Service (FWS), NBEP obtained an inventory of current coastal wetlands, the 500-foot buffer zone, and potential wetland restoration sites for the estuary. While knowing the current state of these resources is vital to managing the resource, an analysis of trends in these resources would help identify threats and put the presentday resources in a historic context.

In 1999, the NBEP and the FWS modified an existing cooperative agreement to produce wetland trends information for the Narragansett Bay Estuary. The FWS works in partnership with UMass (Department of Plant and Soil Sciences, Natural Resources Assessment Group - NRAG) to conduct wetland mapping, trend analysis, and other studies requiring interpretation of aerial photography. NBEP also has an agreement with the URI to perform the geographic information system (GIS) services. URI also played a major role in this project by providing these services. The NBEP will use the results of this work to help develop a coastal wetland conservation and restoration strategy for the Narragansett Bay Estuary.

This report presents the results of this multi-agency cooperative project. It summarizes data for the entire estuary and for several pilot study areas where trends were analyzed back to the 1930s.

## **Study Area**

The Narragansett Bay Estuary is a 147-square mile coastal embayment (including Mount Hope Bay) that dominates the Rhode Island landscape (Figures 1 and 2). It is the receiving basin for seven major watersheds in Rhode Island and Massachusetts including the Blackstone, Moshassuck, Pawtuxet, Taunton, Ten Mile, Warren, and Woonaquatucket. The Estuary is defined by the limits of brackish tidal water and hydrogeomorphology. The baywide coastal wetlands trends analysis (1950s-1990s) was limited to the Rhode Island portion. Within the Narragansett Bay Estuary, six areas were selected as pilot areas to examine wetland trends from the 1930s-1950s in addition to the 1950s-1990s analysis done baywide: 1) Allins Cove, 2) Calf Pasture Point, 3) Jacobs Point, 4) Palmer River, 5) Sachuest Point, and 6) Wesquage Pond (Figure 3).



Figure 1. Location of the Narragansett Bay Estuary and its drainage area; the general boundary of the estuary is the dark gray-shaded area.




Figure 2. Limits of the Narragansett Bay Estuary as defined for this study.





Figure 3. Location of six pilot areas within the Narragansett Bay Estuary.



## Methods

### Data Compilation

Conventional photointerpretation techniques were used to identify trends in coastal wetlands and the 500-buffer around these wetlands. For the Narragansett Bay study area, trends from the 1950s to the 1990s were determined. For the six pilot study areas (Allins Cove, Calf Pasture Point, Jacobs Point, Palmer River, Sachuset Point, and Wesquage Pond), coastal wetland trends were identified for two time periods: the late 1930s/early 1940s-1950s and the 1950s-1990s. Table 1 summarizes the aerial photography used for the study.

Photointerpretation was performed using mirror stereoscopes. Wetlands and deepwater habitats were classified according to "Classification of Wetlands and Deepwater Habitats of the United States" (Cowardin et al. 1979), the national digital data standard for wetland inventory and reporting on wetland trends. For this study, coastal wetlands include Cowardin's marine and estuarine intertidal wetlands - tidal wetlands with measurable traces of ocean-derived salts. Wetland changes to and from nonwetlands were categorized according to the features presented in Table 2. These features represent modifications of the Anderson et al. (1976) national land use/cover classification system. Multiple codes may be assigned to a change in a given wetland. Wetland trends were marked on acetate overlays attached to aerial photographs. Changes in wetlands and deepwater habitats were interpreted using Bausch & Lomb stereo integration scopes. Land use/cover changes in the 500-foot buffer around coastal wetlands were identified using a Bausch & Lomb stereo zoom transfer scope (ZTS) which was also used to match photointerpreted trends data to 1:24,000 frosted mylar maps (prepared by URI). The mylar overlays showing trends were digitized for GIS analysis. The minimum mapping unit for wetland change polygons was 0.25 acre, although smaller polygons of wetland loss were mapped. For more detailed information on methods, see Huber and Nuerminger (2003).

Table 1. Aerial photography used for this study. Note: The 1990s photographs for pilot study areas were the same as used baywide for this period.

Study Area	Study Period	Aerial Photography Used		
		Scale	Emulsion	Date
Entire Bay	1990s	1:40,000	True Color	8/11/96
		1:12,000	True Color	7/6/96
	1950s	1:24,000	Black&White	10-11/51; 5/52
Allins Cove	1930s	1:28,000	Black&White	12/13/38
	1950s	1:20,000	Black&White	5/15/52
Calf Pasture Point	1930s	1:28,000	Black&White	12/13/38
	1950s	1:20,000	Black&White	10/26/51
Jacobs Point	1930s	1:28,000	Black&White	12/13/38
	1950s	1:20,000	Black&White	10/21/51
Palmer River	1930/40s	1:28,000	Black&White	12/13/38; 10/24/41
	1950s	1:20,000	Black&White	10/21/51
Sachuest Point	1930s	1:28,000	Black&White	12/13/38
	1950s	1:20,000	Black&White	10/21/51
Wesquage Pond	1940s	1:28,000	Black&White	10/8/41
	1950s	1:20,000	Black&White	10/26/51

Table 2. Causes of wetland losses, gains, and changes in type.

<b>Cause</b>	<b>Brief Definition</b>
Agriculture	Area subject to farming practices including cropland, orchards, nurseries, vineyards, ornamental horticulture, pasture and hayfields
Barren Land	Nonvegetated or sparsely vegetated lands including mixed, sandy areas (not beaches), strip mines, quarries, and gravel pits
Coastal Processes	Natural processes associated with tidal currents and wave action including erosion, accretion, and dune migration (overwash)
Commercial & Services	Commercial and institutional structures, marinas, paved surfaces, unpaved surfaces, recreational structures, wharves, piers, and shipyards
Ditching	Shallow linear excavation designed to improve drainage; ditches may be filled in to restore wetland hydrology
Erosion from Boat Traffic	Shoreline erosion caused by wakes generated by boats (limited to marina areas)
Excavation	Removal of earth or soil from wetlands or bay and channel bottoms
Forest	Wooded area dominated by trees (deciduous, evergreen, or mixed)
Industrial & Commercial Complexes	Development involving a mixture of factories and business establishments
Jetties & Groins	Artificial rocky structures to maintain navigable channels (jetty) or beaches (groin); these structures may be built or removed
Oyster Colonization	Establishment of an oyster reef
Rangeland	Old fields and thickets (herbaceous, shrub and brush, or mixed cover)
Residential Development	Houses and apartments including lawns
Soil Deposition	Fill material from upland sources deposited in wetlands or waters
Spoil Deposition	Dredged material deposited in wetlands or waters
Tidal Restriction	Tidal flow limited by roadways, railroad embankments, undersized culverts, or similar structures
Transportation, Communications & Utilities	Roads, highways, railroads, powerlines, and similar structures
Unknown	Cause not determined
Urban	Development associated with towns and cities including golf courses and landfills
Vegetation Change	Succession; change in plant composition (specific species noted include <u>Iva frutescens</u> , <u>Phragmites australis</u> , <u>Typha angustifolia</u> )

## Geospatial Database Construction and GIS Analysis

Geospatial database construction was performed by URI's Environmental Data Center (EDC). Each basemap was registered on the digitizing tablet with a RMS value  $\leq 0.003$ . All features delineated for this project were digitized in ArcEdit and coded using ArcGIS 8.2 software. Data for each quad were digitized separately and joined to form one complete baywide coverage. Data for each USGS quadrangle were digitized, coded and proofed before moving on to the next quadrangle. Proofing took place in two phases: 1) on screen in ArcGIS 8.2 to check for coding errors as well as feature errors and 2) a proof plot of the linework information was made and sent along with the mylar basemap for NRAG to proof. Any feature omission or coding change was noted on the proof plot and returned to EDC for final editing.

The land use/cover data were digitized into an existing coverage containing the upland shoreline features from the coastal wetlands data layer and the 500-foot buffer line. Each quad was digitized and proofed separately to be *MAPJOINED* after all land use/cover data were completed. For those polygons coded as freshwater wetland, an item ENHANCED was added and attributed with a Cowardin et al. (1979) classification.

Upon construction of the final digital database, summary tables were generated by EDC using Arc/Info *FREQUENCY* command. These tables were used to prepare tables for this report (in the Results section and Appendices A and B). The database was used to prepare thematic maps showing wetland trends for the estuary and for each pilot area. The maps are presented in a separate folder and hyperlinked to the report.



Palmer River salt marsh (F. Golet photo)



## Results

### Baywide 1996 Status

#### Coastal Wetlands and Waters

In 1996, the Narragansett Bay Estuary (NBE) had 127,742 acres of tidal and subtidal saltwater-influenced habitats (Table 3). The Bay itself (estuarine and marine deepwater habitat) predominates this tidal ecosystem, accounting for 95% of this acreage. Intertidal habitats occupy only 5% of the estuary. Estuarine tidal marshes and swamps comprise 58% of this intertidal habitat, with the remainder made up mostly of nonvegetated tidal unconsolidated shores. The latter includes sandy beaches, sand and mud flats, and cobble-gravel shores. Nine acres of oyster reefs were inventoried.

Over 1,700 acres of vegetated coastal wetlands were altered by ditching and/or impoundment (Table 4). This acreage represented 48% of the NBE's coastal marshes (including estuarine scrub-shrub wetlands). Eighty-eight percent of this acreage was ditched. Only 36 acres of nonvegetated wetlands were altered. Fifteen acres of unconsolidated shore were created by spoil disposal, while nearly 5 acres of rocky shore were created by rip-rap (e.g., groins).

#### 500-Foot Buffer Zone

The 500-foot buffer zone surrounding Narragansett Bay's coastal wetlands accounted for nearly 24,000 acres in 1996 (Table 5). Of this, 35% was represented by residential development (80% single family residences and 18% lawns). Forests and rangeland occupied 22% and 15% of the buffer, respectively. See Table 8 for more detailed findings.



Sachusset Point shoreline (F. Golet photo)



Table 3. 1996 status of coastal wetlands and waters in the Narragansett Bay Estuary (RI).  
(Note: These data summarize totals for mapped polygons only; linear data are not included.) EM=emergent; US=Unconsolidated Shore.

<b>Wetland or Waterbody Type</b>	<b>1990s Acreage</b>
Estuarine Water	
Saline/Brackish	87,764.7
Oligohaline	143.2
-----	-----
Subtotal	87,907.9
Estuarine Marsh	
Emergent Regularly Flooded	242.6
Phragmites Irregularly Flooded	214.6
EM/Phragmites Irregularly Flooded	10.9
EM/US Regularly Flooded	5.8
EM/US Irregularly Flooded	0.3
Emergent Irregularly Flooded	2,185.5
Phragmites/Shrub Irregularly Flooded	3.3
EM/Shrub Irregularly Flooded	6.9
-----	-----
Subtotal	2,659.9
Estuarine Oligohaline Marsh	
Emergent Regularly Flooded	0.8
Phragmites Irregularly Flooded	133.7
EM/Phragmites Irregularly Flooded	113.1
Emergent Irregularly Flooded	108.4
-----	-----
Subtotal	356.04
Estuarine Scrub-Shrub Wetland	
Deciduous Irregularly Flooded	161.8
Shrub/EM Irregularly Flooded	0.7
-----	-----
Subtotal	162.5
Estuarine Reef	
Mollusc (Oyster)	5.9
Estuarine Streambed	
Sand and Mud Regularly Flooded	1.2

Table 3. (continued)

Estuarine Rocky Shore	
Bedrock Regularly Flooded	29.1
Bedrock Irregularly Flooded	96.9
Rubble Regularly Flooded	76.6
Rubble Irregularly Flooded	16.1
-----	-----
Subtotal	218.7
Estuarine Unconsolidated Shore	
Cobble-Gravel Regularly Flooded	68.1
Cobble-Gravel Irregularly Flooded	59.6
Sand Irregularly Exposed	254.4
Sand Regularly Flooded	438.6
Sand/Cobble-Gravel Regularly. Flooded	42.1
Sand/Emergent Regularly Flooded	5.9
Sand Irregularly Flooded	569.4
Mud Irregularly Exposed	198.3
Mud Irregularly. Exposed Oligohaline	0.9
Mud Regularly Flooded	85.5
Mud Regularly. Flooded Oligohaline	7.0
-----	-----
Subtotal	1,729.9
Estuarine Salt Panne	
Irregularly Exposed	39.5
Irregularly Exposed Oligohaline	0.8
Regularly Flooded	1.7
-----	-----
Subtotal	42.0
-----	
<b>Total Estuarine Habitat</b>	<b>93,068.7</b>
-----	
Marine Water	
Unconsolidated Bottom	34,132.9
-----	-----
Subtotal	34,132.9
Marine Rocky Shore	
Regularly Flooded	142.5
Irregularly Flooded	202.2
-----	-----
Subtotal	344.7

Table 3. (continued)

Marine Unconsolidated Shore	
Cobble-Gravel Regularly Flooded	5.9
Cobble-Gravel Irregularly Flooded	9.6
Sand Irregularly Exposed	2.3
Sand Regularly Flooded	100.7
Sand Irregularly Flooded	77.2
-----	-----
Subtotal	195.7
-----	
<b><i>Total Marine Habitats</i></b>	<b><i>34,673.3</i></b>
-----	
<b>Narragansett Bay Grand Total</b>	<b>127,742</b>
-----	

Table 4. Extent of altered coastal wetlands for the Narragansett Bay Estuary in 1996.

<b>Wetland Type</b>	<b>Type of Alteration</b>	<b>Acreage</b>
Emergent		
Regularly Flooded	ditched	0.7
	impounded	6.2
	(subtotal)	(6.9)
Irregularly Flooded	ditched	1336.0
	ditched/impounded	115.2
	impounded	51.7
	(subtotal)	(1502.9)
Emergent Oligohaline		
Regularly Flooded	impounded	0.5
Irregularly Flooded	ditched	19.0
	ditched/impounded	5.6
	impounded	143.7
	(subtotal)	(168.3)
Reef	impounded	3.2
Rocky Shore	artificial	4.7
Scrub-Shrub	ditched	33.9
	ditched/impounded	1.6
	impounded	1.2
	(subtotal)	(36.7)
Unconsolidated Shore	ditched	3.7
	impounded	9.1
	spoil	15.0
	(subtotal)	(27.8)
All Types		1,751.0

Table 5. Land use/cover in the 500-foot buffer around coastal wetlands in the Narragansett Bay Estuary (RI) in 1996. (Note: % buffer totals 103.6% due to round-off procedures.)

<b>Land Use/Cover</b>	<b>Acreage</b>	<b>% of Buffer</b>
Residential	8,737.4	36.9
Commercial	2,179.5	9.2
Industrial	106.1	0.4
Transportation, Communications, Utilities	736.7	3.1
Other Urban or Built-up Land	789.7	3.3
Agriculture	1,318.1	5.6
Rangeland	3,790.6	16
Forest	4,918.8	20.7
Water and Freshwater Wetland	1,542.1	6.5
Barren Land	430.5	1.8
-----	-----	-----
Total	23,706.7	103.6

## Baywide Trends 1951/2 to 1996

### Coastal Wetlands

From the 1950s to the 1990s, the NBE experienced a net loss of 548 acres of tidal habitat. The losses concentrated on intertidal habitats with 306 acres of net loss of estuarine marshes (excluding oligohaline marshes) and a net loss of 205 acres of intertidal nonvegetated wetlands (estuarine unconsolidated shores). During this period, 7.2% of the NBE's estuarine intertidal wetland acreage was lost. Nearly 10% of the estuarine marsh acreage (excluding oligohaline marshes) was lost. Almost 110 acres of coastal waters were lost. Details are provided in Table 6.

The nature and causes of coastal wetland changes are summarized in Table 7. Please note that a loss of a given wetland may be attributed to more than one cause, so the acreage totals from this table may be greater than the net acreage figures reported in Table 6. Causes of wetland changes are illustrated in Figures 4 through 7. Over 50% of the loss of estuarine marsh was due to filling that created upland (dryland) (Figure 4). Nearly 40% of the loss was attributed to conversion to open water (15%), palustrine wetland (12%), and tidal flats (11%). Nine percent of the loss was represented by acreage that changed to estuarine scrub-shrub wetland. While estuarine marshes experienced net losses, there were some gains in estuarine wetland acreage in places. Gains largely came from tidal flats and estuarine water which accounted for over 70% of the estuarine marsh acreage gained (Figure 5). Of the changes to estuarine scrub-shrub wetlands, nearly 60% was due to a gain from estuarine emergent wetland (Figure 6). Forty percent of the changes in these shrub swamps were losses to estuarine marshes (33%) and to upland (7%). Most of the change in estuarine nonvegetated flats and shores were losses (Figure 7). More acreage was converted to open water than came from open water (Table 7). This may be a sign of the impact of rising sea level associated with global warming. About 106 acres of nonvegetated coastal wetlands were converted to upland. (Note: See Appendix A for more detailed findings.)

The locations of these changes are shown on a series of maps. To access information for individual towns, click on the town name: [Barrington](#), [Bristol](#), [Cranston](#), [East Greenwich](#), [East Providence](#), [Jamestown](#), [Little Compton](#), [Middletown](#), [Narragansett](#), [Newport](#), [North Kingstown](#), [Pawtucket](#), [Portsmouth](#), [Providence](#), [South Kingstown](#), [Tiverton](#), [Warren](#), and [Warwick](#).

### 500-foot Buffer Zone Around Coastal Wetlands

Significant changes in the buffer occurred during the 40-year study interval. A 40% increase in residential land occurred largely at the expense of rangeland and agricultural land which decreased by 31% and 55%, respectively (Table 8). This increase was mostly (94%) attributed to a rise in single-family homes along the coastal wetlands, whereas 92% of the loss of agricultural land was from pasture and haylands.

Table 6. Trends in coastal wetlands and waters in the Narragansett Bay Estuary(RI) from the 1950s to the 1990s. (Note: These data summarize totals for mapped polygons only; linear data are not included.) EM=emergent; US=Unconsolidated Shore; Phrag=Phragmites australis.

Wetland or Waterbody Type	1950s Acreage	1990s Acreage	Net Change
Estuarine Water			
Saline/Brackish	87,846.79	87,764.7	-82.1
Oligohaline	170.6	143.2	-27.4
-----	-----	-----	-----
Subtotal	88,017.4	87,907.9	-109.5
Estuarine Marsh			
Emergent Regularly Flooded	280.2	242.6	-37.6
Phragmites Irregularly Flooded	117.1	214.6	+87.5
EM/Phrag Irregularly Flooded	14.9	10.9	-4.0
EM/US Regularly Flooded	7.9	5.8	-2.1
EM/US Irregularly Flooded	0.3	0.3	0
Emergent Irregularly Flooded	2,536.2	2,185.5	-350.7
Phrag/Shrub Irregularly Flooded	3.3	3.3	0
EM/Shrub Irregularly Flooded	5.9	6.9	+1.0
-----	-----	-----	-----
Subtotal	2,965.7	2,659.9	-305.9
Estuarine Oligohaline Marsh			
Emergent Regularly Flooded	3.3	0.8	-2.5
Phragmites Irregularly Flooded	60.4	133.73	+73.3
EM/Phrag Irregularly Flooded	39.2	113.1	+73.9
Emergent Irregularly Flooded	180.4	108.4	-72.0
-----	-----	-----	-----
Subtotal	283.34	356.04	+72.7
Estuarine Reef			
Mollusc (Oyster)	7.3	5.9	-1.4
Estuarine Rocky Shore			
Bedrock Regularly Flooded	29.2	29.1	-0.1
Bedrock Irregularly Flooded	97.1	96.9	-0.2
Rubble Regularly Flooded	76.7	76.6	-0.1
Rubble Irregularly Flooded	15.9	16.1	+0.2
-----	-----	-----	-----
Subtotal	218.9	218.7	-0.2

Table 6. (continued)

Estuarine Streambed			
Sand and Mud Regularly Flooded	0.2	1.2	+1.0
Estuarine Scrub-Shrub Wetland			
Deciduous Irregularly Flooded	128.2	146.4	+18.2
Shrub/EM Irregularly Flooded	0.7	0.7	0
-----	-----	-----	-----
Subtotal	128.9	147.1	+18.2
Estuarine Unconsolidated Shore			
Cobble-Gravel Regularly Flooded	54.7	68.1	+13.4
Cobble-Gravel Irregularly Flooded	55.2	59.6	+4.4
Sand Irregularly Exposed	333.6	254.4	-79.2
Sand Regularly Flooded	440.8	438.6	-2.2
Sand/Cobble-Gravel Reg. Flooded	39.3	42.1	+2.8
Sand/Emergent Regularly Flooded	5.9	5.9	0
Sand/EM Irregularly Flooded	0.5	0	-0.5
Sand Irregularly Flooded	643.5	569.4	-74.1
Sand Reg. Flooded Oligohaline	82.1	0	-82.1
Sand Irreg. Flooded Oligohaline	3.5	0	-3.5
Mud Irregularly Exposed	224.1	198.3	-25.8
Mud Irreg. Exposed Oligohaline	0.9	0.9	0
Mud Regularly Flooded	48.0	85.5	+37.5
Mud Reg. Flooded Oligohaline	2.3	7.0	+4.7
-----	-----	-----	-----
Subtotal	1,934.4	1,729.9	-204.6
Estuarine Salt Panne			
Irregularly Exposed	56.6	39.5	-17.1
Irregularly Exposed Oligohaline	0.8	0.8	0
Regularly Flooded	2.9	1.7	-1.2
-----	-----	-----	-----
Subtotal	60.3	42.0	-18.3
-----			
<b>Total Estuarine Habitat</b>	<b>93,616.5</b>	<b>93,068.7</b>	<b>-547.0</b>
-----			

(Marine totals on following page)



Table 6. (continued)

## Marine Water

Unconsolidated Bottom	34,133.7	34,132.9	-0.8
-----	-----	-----	-----
Subtotal	34,133.7	34,132.9	-0.8
Marine Rocky Shore			
Regularly Flooded	142.8	142.5	-0.3
Irregularly Flooded	201.9	202.2	+0.3
-----	-----	-----	-----
Subtotal	344.7	344.7	0
Marine Unconsolidated Shore			
Cobble-Gravel Regularly Flooded	5.9	5.9	0
Cobble-Gravel Irregularly Flooded	9.6	9.6	0
Sand Irregularly Exposed	2.3	2.3	0
Sand Regularly Flooded	94.9	100.7	+5.8
Sand Irregularly Flooded	83.0	77.2	-5.8
-----	-----	-----	-----
Subtotal	195.7	195.7	0
-----	-----	-----	-----
<b><i>Total Marine Habitats</i></b>	<b><i>34,674.1</i></b>	<b><i>34,673.3</i></b>	<b><i>-0.8</i></b>
-----	-----	-----	-----
<b>Narragansett Bay Grand Total</b>	<b>128,290.4</b>	<b>127,742</b>	<b>-547.8</b>
-----	-----	-----	-----

Table 7. Nature and causes of coastal wetland changes in the Narragansett Bay Estuary from the 1950s to the 1990s. Note: The acreage of areas of change affected by multiple causes has been listed under each of the relevant causes, so acreage totals in this table exceed actual acreage of loss or gain for each coastal wetland type as reported in Table 6.

<b>Wetland Type*</b>	<b>Acreage Affected</b>	<b>Gain From or Lost To</b>	<b>Major Causes (% of affected acreage)</b>
E2EM	52.6	From open water	coastal processes (67), succession (15)
	87.1	From E2US	tidal restriction (48), coastal processes (37)
	33.4	From E2SS	Phragmites invasion (55), ditching (36)
	8.8	From P-wetland	tidal restriction (36), ditching (31), excavation/impoundment (23)
	16.4	From upland	coastal processes (48), unknown (28)
	50.9	To open water	coastal processes (49), tidal restriction (31)
	38.3	To E2US	coastal processes (85)
	0.5	To E2SB	coastal processes (100)
	78.8	To E2SS	<u>Iva</u> succession (61), succession following ditching (33)
	111.1	To P-wetland	ditching (41), tidal restriction (37), succession (11)
	189.8	To upland	rangeland (36), residential (19), commercial/services (14), transportation/utilities (13)
	280.6	Change in EM type	Phragmites (59), other succession (20), tidal restriction (9, excluding Phragmites)
E2SS	0.8	From E2US	coastal processes (100)
	78.8	From E2EM	<u>Iva</u> succession (61), succession/ditching (33)
	33.1	To E2EM	Phragmites (56), succession/ditching (36)
	6.0	To upland	commercial/services (33), forest (27), industrial/commercial (14), agriculture (9), residential (9)

Table 7. (continued)

<b>Wetland Type*</b>	<b>Acreage Affected</b>	<b>Gain From or Lost To</b>	<b>Major Causes (% of affected acreage)</b>
E2US	140.5	From open water	coastal processes (89)
	36.6	From E2EM	coastal processes (89)
	1.5	From E2RS	coastal processes (83)
	34.0	From upland	coastal processes (80)
	250.1	To open water	coastal processes (99)
	112.3	To E2EM	succession (40), tidal restriction (37), coastal processes (17)
	0.8	To E2SS	coastal processes (100)
	21.5	To P-wetland	tidal restriction (52), succession (44)
	105.3	To upland	golf course (33), rangeland (30), barren land (14), commercial/services (5)
	48.5	Change in Type	coastal processes (73)

---

\*E2EM - estuarine emergent wetland; E2SS - estuarine scrub-shrub wetland; E2US - estuarine unconsolidated shore; E2SB - estuarine streambed; E2RS - estuarine rocky shore; P - palustrine.

Figure 4. Percent loss of estuarine emergent wetland in the Narragansett Bay Estuary.




Figure 5. Percent gain in estuarine emergent wetland in the Narragansett Bay Estuary.




Figure 6. Percent change in estuarine scrub-shrub wetland in the Narragansett Bay Estuary.




Figure 7. Percent change in estuarine unconsolidated shore in the Narragansett Bay Estuary.




Table 8. Land use/cover changes (acres and % of 1950s area) in the 500-foot buffer surrounding tidal wetlands in the Narragansett Bay Estuary (RI) from the 1950s to the 1990s.  
+ = gain and - = loss

Land Use/cover Type	50s Acreage	90s Acreage	Acreage Change (% Change)
Residential			
Single-family	4,596.3	6,950.9	+2,354.6 (51)
Lawns	1,474.4	1,561.5	+87.1 (6)
Multi-family	35.3	176.7	+141.4 (400)
Mobile home	13.6	13.8	+0.2 (2)
Other	112.8	34.6	-78.2 (69)
<i>Subtotal</i>	<i>6,232.4</i>	<i>8,737.4</i>	<i>+2,505.0 (40)</i>
Commercial			
Comm.&Institutional Structures	832.3	1,065.5	+233.2 (28)
Wharves, Piers, Shipyards	563.6	558.1	+5.5 (1)
Paved Surfaces	123.9	253.4	+129.5 (100)
Marinas	129.3	201.4	+72.2 (56)
Unpaved Surfaces	90.2	48.3	-41.8 (46)
Recreational Structures	36.9	51.5	+14.6(40)
Junkyard	0.1	0.1	0 (0)
Other	1.2	1.2	0 (0)
<i>Subtotal</i>	<i>1,777.5</i>	<i>2,179.5</i>	<i>+402.1 (23)</i>
Industrial	243.7	90.2	-153.5 (63)
Industrial & Commercial Complexes	23.8	15.9	-7.9 (33)
Transportation, Communications, & Utilities	401.3	736.7	+335.4 (84)
Other Urban or Built-up Land			
Golf Courses	229.8	377	+147.1 (64)
Landfills	18.3	38.8	+20.5 (112)
Cemetaries	48.7	52.5	+3.8 (8)
Other	140.2	321.5	+181.3 (129)
<i>Subtotal</i>	<i>437.0</i>	<i>789.7</i>	<i>+352.7 (81)</i>
Agriculture			
Pasture/hayfields	1,997.2	491.8	-1,505.4 (75)
Cropland	893.00	773.2	-119.8 (13)
Orchards, Nurseries, Vineyards	51.3	49.7	-1.6 (3)
Confined Feeding Lots	6.9	3.4	-3.5 (52)
<i>Subtotal</i>	<i>2,948.4</i>	<i>1,318.1</i>	<i>-1,630.3 (55)</i>

Table 8. (continued)

<b>Land Use/cover Type</b>	<b>50s Acreage</b>	<b>90s Acreage</b>	<b>Acreage Change (% Change)</b>
Rangeland			
Herbaceous	1,073.9	422.2	-651.7 (61)
Shrub and Brush	3,143.4	2,572.6	-570.9 (18)
Mixed	1,301.7	795.9	-505.9 (39)
<i>Subtotal</i>	<i>5,519.1</i>	<i>3,790.6</i>	<i>-1,728.5 (31)</i>
Forest			
Deciduous	2,088.1	2,185.8	+97.7 (5)
Evergreen	234.5	13.6	-220.9 (94)
Mixed	2,145.0	2,719.3	+574.3 (27)
<i>Subtotal</i>	<i>4,467.7</i>	<i>4,918.8</i>	<i>+451.1 (10)</i>
Water & Wetlands			
Vegetated Freshwater Wetland	1,275.9	1,371.3	+95.5 (7)
Nonvegetated Freshwater Wetland	5.00	8.4	+3.5 (69)
Fresh Water	162.2	162.3	+0.1 (0)
<i>Subtotal</i>	<i>1,443.0</i>	<i>1,542.1</i>	<i>+99.0 (7)</i>
Barren Land			
Beaches	18.2	0	-18.2 (100)
Other Sand Areas	188.3	128.9	-59.4 (32)
Mixed Barren Land	277.2	223.5	-53.6 (19)
Strip Mines	10.0	33.4	+23.4 (234)
Bare Exposed Rock	8.6	3.4	-5.2 (61)
Transitional Area	44.6	41.3	-3.3 (7)
<i>Subtotal</i>	<i>546.9</i>	<i>430.5</i>	<i>-116.5 (21)</i>

## Trends for Pilot Study Areas

Wetland trends from the 1930s to the 1950s and the 1950s to the 1990s were examined for six study areas in the Narragansett Bay Estuary: 1) Allins Cove (including West Shore of Barrington), 2) Calf Pasture Point (North Kingstown), 3) Jacobs Point (Warren), 4) Palmer River (Warren), 5) Sachuest Point (Middletown), and 6) Wesquage Pond (Narragansett). All sites experienced net losses of coastal wetlands (Table 9). With a net loss of 104.0 acres, Calf Pasture Point lost the most coastal wetland acreage between the 1930s and the 1990s. Wesquage Pond was next ranked with a net loss of 52.6 acres, followed by Sachuest Point (net loss of 27.9 acres). The other areas experienced only minor net losses (Allins Cove - 7.4 acres; Jacobs Point - 4.4 acres; Palmer River - 0.7 acre). The nature and causes of changes in wetlands and deepwater habitats are presented for each study area in Tables 10 through 15. More detailed findings are given in Appendix B.

The location of these changes are documented on a series of maps showing trends from the 1930s to the 1950s and from the 1950s to the 1990s. To view the maps, [click here](#).

Calf Pasture Point lost more acreage of coastal marsh prior to the 1950s, while it lost more unconsolidated shore (e.g., flats) since then (Table 11). In the earlier period, roughly 70 acres of marsh were lost, with 83% converted to upland; 17 acres of tidal flats were lost with about 14 acres filled (10 acres - commercial/services). Most of this new land was undeveloped in the 1950s (e.g., barren land and rangeland). The rest of the lost marsh was classified as irregularly flooded nonvegetated wetland (spoil deposits in the high marsh) which likely were converted to upland thereafter. From the 50s to the 90s, Calf Pasture Point lost 86 acres of tidal flat and 17 acres of coastal marsh. About 60% of the former losses resulted in an increase in estuarine open water possibly due to a combination of coastal processes (erosion) before the shoreline was stabilized. Filling at Calf Pasture Point created nonvegetated wetlands from open water during the earlier period (this operation was ongoing in the 1950s) and as more fill was deposited these areas were converted to upland. Most of the marsh loss in this area took place during the early stages of this filling operation. By the 1990s, much of the lost coastal marsh between the 1950s and 1990s had become palustrine Phragmites marsh.

Wesquage Pond lost most of its tidal flats prior to the 1950s, accounting for 87% of the losses between the 1930s and 1990s (Table 15). Nearly all of these losses were attributed to tidal restriction which converted intertidal flats mostly to estuarine open water (oligohaline). This action also affected tidal marshes contributing to about a one-acre gain and a five-acre change in tidal marsh type (i.e., some irregularly flooded wetland to regularly flooded marsh and creating oligohaline conditions). About five acres of tidal marshes were filled in Wesquage Pond between the 1950s and the 1990s, with most being undeveloped (rangeland) in the 1990s. About four acres of marsh became open water due to tidal restriction.

Sachuest Point lost most of its coastal wetlands from the 1950s to the 1990s (Table 14). Thirty-eight acres of emergent wetlands were filled during this time. Filling most likely



took place prior to passage of the tidal wetland protection act. Spoil deposition was a major factor impacting wetlands from the 1930s into the 1950s. In the 1990s, much of this acreage remained undeveloped in shrub or herbaceous cover. Some filling also took place at Sachuest Point between the 1930s and 1950s with about 6 acres of tidal flat (estuarine unconsolidated shore) impacted.



High-tide bush marsh at Patience Island (F. Golet photo)

Table 9. Status and trends in coastal wetlands for specific study areas.

Study Area	Wetland Type*	1930s Acreage	1950s Acreage	Net Change in Acreage (% Change)	1990s Acreage	Net Change in Acreage (% Change)	Total Change 1930s-1990s (% Change)
Allins Cove	EEM	65.8	62.7	-3.2 (-5)	45.7	-17.0 (-27)	-20.1 (-31)
	EEMO	13.7	7.2	-6.5 (-47)	8.7	+1.5 (+21)	-5.0 (-37)
	ESS	1.1	0.4	-0.7 (-64)	5.9	+5.5 (+1375)	+4.8 (+436)
	EUS	22.3	20.2	-2.1 (-9)	35.2	+15.0 (+74)	+12.9 (+58)
Calf Pasture Point	EEM	128.1	66.8	-61.3 (-48)	50.1	-16.7 (-25)	-78.0 (-61)
	EEMO	18.5	18.8	+0.3 (+2)	14.9	-3.9 (-21)	-3.6 (-20)
	ESS	5.3	0	-5.3 (-100)	4.4	+4.4 (NA)	-0.9 (-17)
	EUS	42.5	100.0	+57.5 (+135)	20.8	-79.2 (-79)	-21.7(-51)
	ERS	0.3	0.3	0	0.5	+0.2 (+67)	+0.2 (+67)
Jacobs Point	EEM	22.3	22.3	0	23.9	+1.6 (+7)	+1.6 (+7)
	EEMO	9.7	7.1	-2.6 (-27)	12.6	+5.5 (+78)	+2.9 (+30)
	ESS	12.7	12.7	0	1.8	-10.9 (-86)	-10.9 (-86)
	EUS	7.3	7.3	0	9.3	+2.0 (+27)	+2.0 (+27)
	ERS	0.7	0.7	0	0.7	0	0
Palmer River	EEM	214.9	212.7	-2.2 (-1)	219.3	+6.6 (+3)	+4.4 (+2)
	EEMO	1.2	0	-1.2 (-100)	0	0	-1.2 (100)
	ESS	15.2	15.2	0	9.0	-6.2 (-41)	-6.2 (-41)
	EUS	8.1	8.7	+0.6 (+8)	10.4	+1.7 (+20)	+2.3 (+28)

\* EEM - estuarine emergent; EEMO - estuarine emergent oligohaline; ESS - estuarine scrub-shrub; EUS - estuarine unconsolidated shore; ERS - estuarine rocky shore; ESB - estuarine streambed; MUS - marine unconsolidated shore; MRS - marine rocky shore.

Table 9. (continued)

Study Area	Wetland Type	1930s Acreage	1950s Acreage	Net Change in Acreage (% Change)	1990s Acreage	Net Change in Acreage (% Change)	Total Change 1930s-1990s (% Change)
Sachuest Point	EEM	62.6	69.9	+7.3 (+12)	32.2	-37.7 (-54)	-30.4 (-49)
	EEMO	3.1	1.9	-1.2 (-39)	14.3	+12.4 (+653)	+11.2 (+361)
	ESS	5.3	0	-5.3 (-100)	0	0	-5.3 (-100)
	EUS	20.6	14.4	-6.2 (-30)	17.2	+2.8 (+19)	-3.4 (-17)
	ERS	2.0	2.0	0	2.0	0	0
	MUS	46.7	46.7	0	46.7	0	0
	MRS	34.4	34.4	0	34.4	0	0
Wesquage Pond	EEM	7.7	8.4	+0.7 (+9)	0	-8.4 (-100)	-7.7 (-100)
	EEMO	19.3	19.6	+0.3 (+2)	24.1	+4.5 (+23)	+4.8 (+25)
	ESS	0.4	0.3	-0.1 (-25)	0	-0.3 (-100)	-0.4 (-100)
	EUS	51.4	2.3	-49.1 (-96)	1.7	-0.6 (-26)	-49.7 (-97)
	EUS/EM	0.2	0.5	+0.3 (+150)	0	-0.5 (-100)	-0.2 (-100)
	ESB	0.2	0.2	0	0.6	+0.4 (+200)	+0.4 (+200)
	ERS	11.5	11.5	0	11.7	+0.2 (+2)	+0.2 (+2)
	MUS	11.8	11.8	0	11.8	0	0
	MRS	3.5	3.5	0	3.5	0	0

Table 10. Nature and causes of coastal wetland and deepwater habitat trends for Allins Cove.

Time Period	Wetland Type*	Change Type	Acreage	Causes
1930s-50s	NVW	loss	2.9	coastal processes, filling (residential development)
		gain	0.7	coastal processes
		type change	1.0	coastal processes
		no change	18.5	n/a
	VW	loss	11.0	coastal processes, filling (golf course)
		gain	0.6	coastal processes, unknown
		type change	0.7	<u>Phragmites</u> , unknown
		no change	69.0	n/a
	CW	loss	1.0	coastal processes, unknown
		gain	5.4	coastal processes
		no change	20.7	n/a
		no change	20.7	n/a
1950s-90s	NVW	loss	1.0	coastal processes
		gain	15.9	coastal processes, unknown
		type change	2.2	coastal processes
		no change	17.0	n/a
	VW	loss	14.0	tidal restriction, filling (golf course, residential development), coastal processes
		gain	4.1	coastal processes, <u>Phragmites</u> invasion
		type change	6.9	ditching/ <u>Iva</u> succession, tidal restriction, <u>Phragmites</u> , unknown
		no change	46.2	n/a
	CW	loss	16.2	coastal processes, spoil deposition, <u>Phragmites</u> invasion
		no change	9.9	n/a
		no change	9.9	n/a
		no change	9.9	n/a

\* NVW - nonvegetated wetland; VW - vegetated wetland; CW - coastal water (deepwater habitat); n/a - not applicable.

Table 11. Nature and causes of coastal wetland and deepwater habitat trends for Calf Pasture Point.

Time Period	Wetland Type*	Change Type	Acreage	Causes
1930s-50s	NVW	loss	17.1	filling (commercial/services, barren land), coastal processes
		gain	74.5	spoil deposition, coastal processes
		type change	0.6	spoil deposition
		no change	25.0	n/a
	VW	loss	70.3	filling (barren land, rangeland, commercial/services, spoil deposition), coastal processes, ditching/succession,
		gain	4.0	coastal processes, <u>Phragmites</u> , unknown
		type change	8.4	spoil deposition, unknown
		no change	73.2	n/a
	CW	loss	123.0	filling (spoil deposition, barren land, commercial/services), coastal processes, <u>Phragmites</u> invasion
		gain	6.1	coastal processes, tidal restriction
		no change	6.0	n/a
		type change	3.7	spoil deposition, coastal processes, jetty/groin removal
1950s-90s	NVW	loss	85.7	coastal processes, filling (rangeland)
		gain	6.2	coastal processes, spoil deposition
		type change	10.9	n/a
		no change	40.1	n/a
	VW	loss	17.2	<u>Phragmites</u> invasion, filling (forest, rangeland, landfill, golf course, spoil deposition), coastal processes, tidal restriction
		gain	8.5	coastal processes, succession/ditching
		type change	28.3	succession/ditching, <u>Phragmites</u> , <u>Iva</u> , spoil deposition, unknown
		no change	5.3	n/a
	CW	loss	5.1	coastal processes, excavation
		gain	54.8	coastal processes
		type change	5.3	n/a
		no change	5.3	n/a

\*NVW - nonvegetated wetland; VW - vegetated wetland; CW - coastal water (deepwater habitat); n/a - not applicable.

Table 12. Nature and causes of coastal wetland and deepwater habitat trends for Jacobs Point.

Time Period	Wetland Type*	Change Type	Acreage	Causes
1930s-50s	NVW	no change	8.0	n/a
	VW	loss	2.6	agriculture, tidal restriction/agriculture
		no change	29.4	n/a
	CW	no change	0.6	n/a
1950s-90s	NVW	loss	1.4	coastal processes
		gain	0.6	coastal processes
		no change	8.0	n/a
	VW	loss	3.8	filling (rangeland, residential development), coastal processes,
		type change	14.5	succession/ditching, <u>Phragmites</u> , <u>Iva</u>
		no change	23.8	n/a
	CW	loss	0.6	coastal processes

\*NVW - nonvegetated wetland; VW - vegetated wetland; CW - coastal water (deepwater habitat); n/a - not applicable.

Table 13. Nature and causes of coastal wetland and deepwater habitat trends for Palmer River.

Time Period	Wetland Type*	Change Type	Acreage	Causes
1930s-50s	NVW	gain	0.6	unknown
		no change	8.1	n/a
	VW	loss	3.4	tidal restriction, filling (commercial/services, barren land, residential), coastal processes
		no change	227.9	n/a
	CW	loss	39.7	impoundment
		gain	0.5	coastal processes
1950s-90s	NVW	no change	10.0	n/a
		gain	2.2	coastal processes, unknown
		loss	0.6	unknown
	VW	no change	8.1	n/a
		gain	3.8	coastal processes, spoil deposition, succession/ditching, unknown
		loss	3.3	filling (residential development, commercial/services, rangeland)
		type change	21.0	<u>Phragmites</u> , succession/ditching, <u>Iva</u> , unknown
	CW	no change	203.6	n/a
		loss	8.0	filling (rangeland, commercial/services, residential development), coastal processes, succession/ditching, unknown

\* NVW - nonvegetated wetland; VW - vegetated wetland; CW - coastal water (deepwater habitat); n/a - not applicable.

Table 14. Nature and causes of coastal wetland and deepwater habitat trends for Sachuest Point.

Time Period	Wetland Type*	Change Type	Acreage	Causes
1930s-50s	NVW	loss	6.2	filling (spoil deposition, commercial/services), coastal processes, <u>Phragmites</u> invasion
		no change	97.5	n/a
	VW	loss	4.8	filling (residential, transportation/comm./utilities, commercial/services)
		gain	5.7	spoil deposition, <u>Phragmites</u> invasion, coastal processes
		type change	29.0	spoil deposition
1950s-90s	CW	no change	37.2	n/a
		no change	2.2	n/a
	NVW	gain	2.8	coastal processes
		type change	1.9	coastal processes
		no change	95.5	n/a
	VW	loss	26.3	filling (spoil deposition, rangeland, commercial/services, barren land)
		gain	1.0	revegetation (sediment accretion after excavation)
		type change	20.6	tidal restriction, <u>Phragmites</u> invasion, succession/ditching
	CW	no change	24.9	n/a
		loss	1.6	revegetation (excavation), coastal processes
		no change	0.6	n/a

\*NVW - nonvegetated wetland; VW - vegetated wetland; CW - coastal water (deepwater habitat); n/a - not applicable.



Table 15. Nature and causes of coastal wetland and deepwater habitat trends for Wesquage Pond.

Time Period	Wetland Type*	Change Type	Acreage	Causes
1930s-50s	NVW	loss	49.1	tidal restriction, coastal processes
		gain	0.3	coastal processes
		no change	29.5	n/a
	VW	loss	0.4	filled (commercial)
		gain	1.5	tidal restriction, coastal processes
		type change	5.0	tidal restriction, <u>Phragmites</u>
		no change	21.9	n/a
	CW	loss	0.5	coastal processes, tidal restriction
		gain	47.6	tidal restriction
1950s-90s	NVW	no change	20.5	n/a
		loss	1.1	filling (residential development)
		gain	0.7	coastal processes, jetty/groin construction
	VW	no change	28.7	n/a
		loss	9.2	tidal restriction, filling (commercial/services, rangeland, residential)
		gain	5.0	tidal restriction, <u>Phragmites</u> , unknown
		type change	4.0	tidal restriction, <u>Phragmites</u> , unknown
	CW	no change	15.0	n/a
		loss	5.7	tidal restriction, <u>Phragmites</u> , filling (residential), jetty/groin construction, unknown
		gain	4.4	tidal restriction
		type change	0.3	impounded/tidal restriction
		no change	62.1	n/a

\* NVW - nonvegetated wetland; VW - vegetated wetland; CW - coastal water (deepwater habitat); na - not applicable.

## Conclusions

The Narragansett Bay Estuary (NBE) contains about 130,000 acres of tidal and subtidal habitats. Open water is the predominant feature of the Bay occupying about 95% of the tidal ecosystem. Intertidal habitats (marshes, beaches, flats, and other shores) represent only 5% of the ecosystem. Of this, vegetated wetlands (mostly salt marshes) comprise 58% of the acreage, with the rest made up mostly of tidal flats. Nine acres of oyster reefs were inventoried. Over 1,700 acres (or 48%) of the coastal marshes have been ditched and/or impounded. Slightly more than one-third of the 500-foot buffer around the coastal wetlands is occupied by residential development. Forests and rangeland (i.e., fields and shrub thickets) represent 21% and 16% of the buffer, respectively.

Between the 1950s and 1990s, the NBE lost a net total of about 110 acres of estuarine open water, nearly 306 acres of salt and brackish marshes, and 205 acres of intertidal shores. A net gain of 73 acres of slightly brackish marshes took place, mostly at the expense of more saline wetlands. About 190 acres of salt/brackish marshes were filled. Common reed (Phragmites australis), a widespread invasive grass, increased its distribution during the study period by roughly 240 acres. Major causes of coastal marsh loss and degradation were filling and tidal restriction. Gains and losses of coastal marsh attributed to coastal processes (erosion/accretion) were nearly even, where these processes caused about 1.5 times more loss of unconsolidated shores than gains between the 1950s and 1990s.

For six areas in the NBE, wetlands trends were examined back to the 1930s (Allins Cove, Calf Pasture Point, Jacobs Point, Palmer River, Sachuest Point, and Wesquage Pond). All sites experienced net losses of coastal wetlands, but only Calf Pasture Point (104 acres), Wesquage Pond (53 acres), and Sachuest Point (28 acres) lost more than 10 acres. The other areas lost less than eight acres each.

## Acknowledgments

Funding for this project was provided by the Narragansett Bay Estuary Program (NBEP). Helen Cottrell served as project officer for NBEP. She reviewed the draft manuscript and products and provided photographs for use in this report. Ralph Tiner was project officer for the U.S Fish and Wildlife Service (FWS). He was responsible for study design, project oversight, data analysis and synthesis, and report preparation.

Wetland trends data were collected by the Natural Resources Assessment Group (NRAG) in the Department of Plant and Soil Sciences, University of Massachusetts-Amherst, under the direction of Dr. Peter Veneman, principal investigator. Most of the photointerpretation and cartographic work was performed by Irene Huber and Todd Nuerminger. Mary Johnson (NRAG) assisted in photointerpretation of land use/cover trends in the 500-foot buffer around coastal wetlands. Craig Polzen also did some work on this photo-analysis and provided GIS support to NRAG.

Digital database construction and GIS analyses were accomplished by Aimée Mandeville of the University of Rhode Island's Environmental Data Center. She digitized map overlay products to create the geospatial database, performed analytical inquiries, and produced color-coded maps and statistical summaries presented in this report including the Appendices. Figure 1 was prepared by Paul Jordan, Rhode Island Department of Environmental Management, Geographic Information System Program, while Aimée created Figures 2 and 3. Herbert Bergquist (FWS) assisted in preparing the cover of this report, while Lorraine Fox (FWS) scanned the photos for conversion to digital images. Dr. Frank Golet, University of Rhode Island, graciously provided color photographs of coastal wetlands for use in this report.

.

## References

Anderson, J.R., E.E. Hardy, J.T. Roach, and R.E. Witmer. 1976. A Land Use and Land Cover Classification System for Use with Remote Sensor Data. U.S. Geological Survey Professional Paper 96A, U.S. Government Printing Office, Washington, DC.

Cowardin, L.W., V. Carter, F.C. Golet, and E.T. LaRoe. 1979. Classification of Wetlands and Deepwater Habitats of the United States. U.S. Fish and Wildlife Service, Washington, DC.

Huber, I. and T. Nuerminger. 2003. Rhode Island Narragansett Bay Project Area: Trends Analysis Methodology. Department of Plant and Soil Sciences, Natural Resources Assessment Group, University of Massachusetts, Amherst, MA.

## **Appendices**

[Click here for Appendices](#)